



Endolysin CHAP- domain

Datasheet, Version 2/2016

protean
Forefront of Biotechnology

Catalog #	1519
Synonyms	Endolysin Φ 812 catalytic domain
Type	Recombinant
Source	E. coli
Species	Bacteriophage Φ 812
Tag	His6
Form	Liquid

Purity	>96% by SDS-PAGE
Shipping	Ice pack

Introduction

Endolysins are hydrolytic enzymes produced by bacteriophages in order to cleave the host's cell wall during the final stage of the lytic cycle. They are highly evolved enzymes able to cleave peptidoglycan (murein), the main component of bacterial cell wall, which allows the release of progeny virions from the lysed cell. They are highly species specific, leaving non-target bacteria intact.

Description

Recombinant CHAP fragment of phage Φ 812 catalytic domain

Application

Anti-Staphylococcus aureus activities, Turbidity-reduction assay, plaque assay, various anti-pathogen studies

Purification method

Affinity chromatography

Formulation

50mM Tris pH 7,5; 150 mM NaCl, 1 mM EDTA; 5 mM DTT, 50% glycerol, may contain traces of imidazol

Specificity

Specifically cleave peptidoglycan in cell wall leading to cell lysis.

Storage

-80C, aliquot to avoid repeated freezing and thawing.

Analyte specific reagent (ASR) manufactured under ISO 13485.

Country of origin: Czech Republic

Protean s.r.o.
Pod Lesem 9
373 16 Dobra Voda u CB
The Czech Republic

ISO 9001 & ISO 13485
CERTIFIED



www.protean.bio
info@protean.bio
+420 380 120 427